



UNIVERSITI PUTRA MALAYSIA

**A STUDY ON THE POTENTIAL OF USING SEAWEED AS
BIOMONITORING INDICATOR IN KISH ISLAND, IRAN**

ALI DADOLAH-SOHRAB

FSAS 2003 45

**A STUDY ON THE POTENTIAL OF USING SEAWEED AS
BIOMONITORING INDICATOR IN KISH ISLAND, IRAN**

By

ALI DADOLAH-SOHRAB

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

August 2003



DEDICATION

To

**Memory of my parents and my sister Maryam whom their spirits will always
be a part of mine**

**My wife Mojgan for years of love and dedication, and
my daughter Paradise whom her presence enriched my life**

and

My sisters whom taught me how to work hard and be happy

Thanks to Allah

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirements for the degree of Doctor of Philosophy

**A STUDY ON THE POTENTIAL OF USING SEAWEED AS
BIOMONITORING INDICATOR IN KISH ISLAND, IRAN**

By

ALI DADOLAH-SOHRAB

August 2003

Chairman: Misri Kusnan, Ph.D.

Faculty: Science and Environmental Studies

The study was carried out to evaluate the use of seaweed as biomonitoring in Kish Island, Iran. Seaweed, sediment and seawater samples were collected bimonthly from June 1999 to April 2000 at 5 different sites. Seaweed samples were collected, identified and biomass recorded for 4 seasons from April 1999 to March 2000. Diversity and biomass differed within sites and seasons. The highest seaweed biomass were recorded from northwest and west sides of study area.

The status of selected heavy metals (Cd, Cu, Ni, Pb, V and Zn) were determined in seawater, sediment and 10 dominant seaweed species. Metals level in seawater ranged from 0.02-0.10, 0.09-1.22, 0.11-0.18, 0.27-2.04 and 0.17-0.27 $\mu\text{g l}^{-1}$ for Cd, Cu, Ni, Pb and Zn, respectively, the metals concentrations were higher during cold seasons compared to the warm seasons. Metal concentrations in sediments ranged from 0.18-0.39, 2.39-4.55, 2.30-11.92, 2.92-5.92, 1.69-4.76, 4.04-9.51 $\mu\text{g g}^{-1}$ dry weight for Cd, Cu, Ni, Pb, V and Zn, respectively. Higher variations in metal concentrations were observed at sites Symorgh and Darakht-e-Sabz compared to other sites.

Metal levels in seaweeds showed considerable variations ranging from 0.44-1.74, 0.76-3.42, 0.37-2.23, 0.96-4.47, 0.53-2.16 and 3.44-10.23 $\mu\text{g g}^{-1}$ dry weight for Cd, Cu, Ni, Pb, V and Zn, respectively. Cadmium was lower in Chlorophyta than the other groups. Chlorophyta accumulated more Zn, V, Pb, Ni and Cu than other groups. In addition, variations in metals contents between species were obvious from the same habitat and in several cases within the same taxonomic groups. Generally, heavy metals level in this study were relatively lower than the other parts of the world and in some cases lower than the other parts of the Persian Gulf.

Toxicity test experiments were carried out using *Cystoseira myrica* from Kish Island, Iran and *Sargassum ilicifolium* species from Port Dickson, Malaysia. All selected concentrations of Cd and V and Cu concentrations at above 0.01 mg l^{-1} showed toxic effects to both species. The toxic effects of Ni and Zn on the *C. myrica* and *S. ilicifolium* were at higher concentrations than the other metals and in addition Cu and V appeared to be more toxic than the other elements.

Using both species bioconcentration factors of Pb, Zn and Cu were higher than the other metals. This could be related to their electronegativity values. Elements uptake by different parts of *S. ilicifolium* during light period decreased in the following order: fronds > receptacles > stipes, whereas during dark period, accumulation of Cd, Cu, V and Ni exhibited similar trend to the above order, but Zn and Pb were higher in receptacles followed by fronds and stipes.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

KAJIAN MENGGUNAKAN RUMPAIRLAUT SEBAGAI PENUNJUK BIOPEMONITORAN DI PULAU KISH, IRAN

Oleh

ALI DADOLAH-SOHRAB

Ogos 2003

Pengerusi: Misri Kusnan, Ph.D.

Fakulti: Sains dan Pengajian Alam Sekitar

Kajian telah dijalankan untuk menilai kegunaan rumput laut sebagai penunjuk pencemaran di Pulau Kish, Iran. Sampel rumput laut, sedimen dan air laut telah diambil setiap 2 minggu dari Jun 1999 hingga April 2000 dari 5 kawasan berlainan. Sampel rumput laut diambil, dikenal pasti dan biomasnya direkodkan untuk 4 musim dari April 1999 hingga March 2000. Diversiti dan biomas berbeza di antara kawasan dan musim. Biomas rumput laut tertinggi telah direkodkan dari kawasan Barat laut dan Barat kawasan kajian.

Status logam berat pilihan (Cd, Cu, Ni, Pb, V dan Zn) telah ditentukan pada air laut, sedimen dan 10 spesis rumput laut utama. Paras logam di dalam air laut berada di antara 0.02-0.10, 0.09-1.22, 0.11-0.18, 0.27-2.04 dan 0.17-0.27 $\mu\text{g l}^{-1}$ untuk Cd, Cu, Ni, Pb dan Zn, logam-logam berada pada paras tertinggi semasa musim sejuk berbanding semasa musim panas. Kepekatan logam dalam sedimen berada di antara 0.18-0.39, 2.39-4.55, 2.30-11.92, 2.92-5.92, 1.69-4.76, 4.04-9.51 $\mu\text{g g}^{-1}$ berat kering untuk Cd, Cu, Ni, Pb, V dan Zn. Variasi kepekatan logam yang tinggi diperolehi dari kawasan Symorgh dan Darakht-e-Sabz berbanding dengan kawasan-kawasan lain. Paras logam dalam rumput laut menunjukkan variasi antara 0.44-1.74, 0.76-

3.42, 0.37-2.23, 0.96-4.47, 0.53-2.16, dan 3.44-10.23 $\mu\text{g g}^{-1}$ berat kering untuk Cd, Cu, Ni, Pb, V dan Zn. Kadmium adalah rendah dalam Chlorophyta berbanding kumpulan-kumpulan lain. Chlorophyta mengumpul Zn, V, Pb, Ni dan Cu yang lebih tinggi berbanding dengan kumpulan-kumpulan lain. Selain itu, variasi dalam kandungan logam-logam antara spesis-spesis adalah ketara sekiranya berasal dari habitat yang sama dan dalam sesetengah kes, dari kumpulan taksonomi yang sama. Secara umumnya, paras logam berat dalam kajian ini adalah rendah berbanding kawasan-kawasan lain di dunia dan dalam sesetengah kes ianya adalah rendah berbanding kawasan lain di Telak Parsi.

Ujian ketoksikan telah dilakukan dengan menggunakan spesis *Cystoseira myrica* dari Pulau Kish, Iran dan *Sargassum ilicifolium* dari Port Dickson, Malaysia. Kesemua kepekatan pilihan untuk Cd, V dan Cu kepekatan melebihi 0.01 mg l^{-1} adalah toksik kepada kedua-dua spesis. Kesan toksik untuk Ni dan Zn terhadap *C. myrica* dan *S. ilicifolium* adalah pada kepekatan yang tinggi berbanding dengan logam-logam lain, dan umumnya Cu dan V terbukti lebih toksik berbanding elemen lain.

Faktor biokepekatan bagi Pb, Zn dan Cu adalah tinggi berbanding logam lain. Ini mungkin ada perkaitan dengan nilai elektronegativiti mereka. Pengambilan elemen oleh bahagian yang berlainan bagi *S. ilicifolium* semasa waktu siang berkurangan mengikut turutan berikut: daun (fronds) > organ seks (receptacles) > batang (stipes), manakala sewaktu malam, pengambilan Cd, Cu, V dan Ni menunjukkan bentuk yang sama seperti di atas, tetapi Zn dan Pb adalah tinggi di organ seks, diikuti dengan daun dan batang.

ACKNOWLEDGEMENTS

I would like to extend my gratitude to the members of my supervisory committee; Dr. Misri Kusnan, Assoc. Prof. Dr. Ahmad Ismail, Dr. Hishamuddin Omar and Assoc. Prof. Dr. Ahmad Savari for their advice, kind and constant support, encouragement and help in all my difficulties.

This research was made possible through the financial support of the AKF (Agha Khan Foundation) which is highly appreciated. I would like to acknowledge support from the Kish University and KFZO (Kish Free Zone Organization) make this study possible.

Furthermore, my special thanks to Mrs. Catherine Hieronymi (AKF Scholarship Officer) and I would like to thank Dr. Askari, Dr. Sam, Dr. Rostami and my colleagues at Kish University, also Mr. Hamid Rezaei, Ibrahim Jalili and Majeed Ismail Kashi, Abu Hena, Nik, Syaizwan, Hidir and all my friends.

Finally, my sincere thanks are to my wife and daughter. I would like also to express my greatest gratitude and thanks to my sisters and their husbands and my parents in law for their encouragement during my study.

I certify that an Examination Committee met on 1st August 2003 to conduct the final examination of Ali Dadolahi-Sohrab on his Doctor of Philosophy thesis entitled “A Study on the Potential of Using Seaweed as Biomonitoring Indicator in Kish Island, Iran” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

ABDUL RAHIM ISMAIL, Ph.D.

Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Chairman)

MISRI KUSNAN, Ph.D.

Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Member)

AHMAD ISMAIL, Ph.D.

Associate Professor
Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Member)

HISHAMUDDIN OMAR, Ph.D.

Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Member)

AHMAD SAVARI, Ph.D.

Associate Professor
Faculty of Oceanography
University of Shaheed Chamran, Ahwaz, Iran
(Member)

NOOR AZHAR MOHAMED SHAZILI, Ph.D.

Professor
Faculty of Science and Technology
College University Science and Technology Malaysia
(Independent Examiner)



GULAM RASUL RAHMAT ALI, Ph.D.

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 29 SEP 2003

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

MISRI KUSNAN, Ph.D.

Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Chairman)

AHMAD ISMAIL, Ph.D.

Associate Professor
Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Member)

HISHAMUDDIN OMAR, Ph.D.

Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Member)

AHMAD SAVARI, Ph.D.

Associate Professor
Faculty of Oceanography
University of Shaheed Chamran, Ahwaz, Iran
(Member)



AINI IDERIS, Ph.D.
Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 26 SEP 2003

DECLARATION

I hereby declare that the thesis based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.


ALI DADOLAH-SOHRAB

Date: 4/11/2003

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xiii
LIST OF FIGURES	xviii
LIST OF PLATES	xxi
LIST OF ABBREVIATIONS	xxiii
 CHAPTER	
 I INTRODUCTION	 1
 II LITERATURE REVIEW	 13
Seaweed	13
Classification of Seaweeds	14
Seaweeds Diversity in the Persian Gulf	18
Distribution and Diversity of Seaweeds	19
Factors Affecting on Distribution and Diversity of Seaweed	20
Heavy Metals	29
Heavy Metals Accumulation by Seaweeds	31
Characteristics of Seaweed as a Biological Indicator	50
Toxicity Test Studies Using Seaweeds	53
Basics of Toxicity Test Technique Using Seaweeds	55
Seaweed Morphology and Anatomy	58
 III METHODOLOGY	 60
Study Area	60
Sampling Sites	60
Sampling Methods	66
Biomass Study	66
Heavy Metals Study in Kish Island	67
Preparation of Samples for Analysis	69
Chemicals	69
Apparatus	70
Cleaning and Drying of Samples	70
Physicochemical Data	72
Seawater Nutrients	72
Bioassay Studies	73
Bioassay Study Using <i>Cystoseira myrica</i>	74



Bioassay Study Using <i>Sargassum ilicifolium</i>	75
End Point Measurement in Seaweeds	76
Sampling of Seaweed and Seawater	77
Photosynthesis and Respiration	77
Chlorophyll Contents	78
Metals Uptake by Different Parts of Seaweed	79
Qualitative Assurance Procedures	81
Statistical Analysis	82
Biomass Study	83
Heavy Metal in Marine Environment	84
Bioassay Study	85
IV RESULTS AND DISCUSSION	87
Seaweed in Kish Island	87
Phylum Chlorophyta	87
Phylum Phaeophyta	92
Phylum Rhodophyta	96
The Ecology of Seaweed in Kish Island	101
Biomass Study	104
Overall Observation about Seaweed Ecology	120
Heavy Metals in Kish Island	127
Heavy Metals in Seawater of Kish Island	127
Heavy Metals in Sediments of Kish Island	134
Heavy Metals in Seaweeds of Kish Island	139
The Use of Seaweed as Biomonitors of Metals	163
Comparison of the Metal Contents in Different Parts of Seaweed	167
Seasonal Changes of Heavy Metals in Seaweeds	170
Metal Pollution Index (MPI)	180
Toxicity Test	183
Toxicity Test on <i>Cystoseira myrica</i> (Kish Island)	183
Toxicity Test on <i>Sargassum ilicifolium</i> (Port Dickson)	195
Overall Observation about Metals Effects on Seaweeds	208
Metals Accumulation in Toxicity Test	215
V GENERAL DISCUSSION AND CONCLUSION	230
BIBLIOGRAPHY	261
APPENDICES	291
BIODATA OF THE AUTHOR	310

LIST OF TABLES

Table		Page
1	Photosynthesis pigments common to divisions and classes of algae.	17
2	Research on the seaweed in the Persian Gulf.	18
3	Essential elements to the growth of seaweed.	27
4	Classification of elements according to toxicity and availability.	30
5	Compilation of published studies of heavy metals content ($\mu\text{g g}^{-1}$ dry weight) of selected tropical and subtroical seaweed species.	34
6	Description of sampling sites.	62
7	Floristic affinity as determined by Cheney's (R+C)/P ratio (Cheney 1977) for published floras along the Persian Gulf Coasts.	103
8	Biomass (g wet wt m^{-2}) of seaweeds at all sites during spring in Kish Island.	105
9	Biomass (g wet wt m^{-2}) of seaweeds at all sites during summer in Kish Island.	106
10	Biomass (g wet wt m^{-2}) of seaweeds at all sites during autumn in Kish Island.	107
11	Biomass (g wet wt m^{-2}) of seaweeds at all sites during winter in Kish Island.	107
12	Average biomass (g wet wt m^{-2}) and per cent values of each class in each station and in each season. Seasonal abundance and annual means are also reported.	109
13	Average annual biomass (g wet wt m^{-2}) of seaweeds at different sites in Kish Island.	111
14	Average biomass (g wet wt m^{-2}) of seaweeds in different season in Kish Island.	112
15	Biomass (g wet wt m^{-2}) of each species in different zones in Kish Island.	113
16	Results of two-way ANOVA based on the biomass of different Classes of seaweeds during different seasons.	121
17	Results of correlation coefficient between total seaweeds biomass with environment parameters.	121



18	The metal concentrations (means \pm S.D.) in $\mu\text{g l}^{-1}$ from the seawater around the Kish Island.	127
19	Results of Kruskal-Wallis test based on estimates of heavy metal concentrations in different locations of Kish Island.	129
20	Results of Kruskal-Wallis test based on estimates of heavy metal concentrations in different seasons of Kish Island.	130
21	Heavy metals concentrations ($\mu\text{g l}^{-1}$) in coastal waters of Kish Island with those from other regions of the world.	133
22	Heavy metals concentrations (means \pm S.D.) in $\mu\text{g g}^{-1}$ dry wt in sediments of Kish Island.	134
23	Results of Kruskal-Wallis test based on estimates of heavy metal concentrations in sediments of Kish Island.	135
24	Background levels of heavy metals in sediments of Persian Gulf.	136
25	Heavy metals concentrations ($\mu\text{g g}^{-1}$ dry wt) in surface sediments from the different parts of the world.	139
26	Correlation between concentrations of different metals in sediment and seawater.	139
27	Range of metal concentrations ($\mu\text{g g}^{-1}$ dry weight) in all species from Kish Island.	140
28	Concentrations (means \pm S.D.) in $\mu\text{g g}^{-1}$ dry wt of heavy meals in three classes of seaweeds around the Kish Island.	141
29	Concentrations (means \pm S.D.) of heavy metals ($\mu\text{g g}^{-1}$ dry weight) of the seaweeds in Kish Island.	142
30	Metal concentrations (means \pm S.D.) in $\mu\text{g g}^{-1}$ dry weight of algae along the coasts of Kish Island.	143
31	Statistical analysis of the metal contents data (two-way of ANOVA).	143
32	List of green algae sampled from different sites.	144
33	Results obtained from the multiple regression analysis of the metal contents in Chlorophyta species in relationship with the metal contamination in sediment and seawater.	144

34	Correlation coefficient between metal concentrations in ten dominant species with sediment and seawater around the Kish Island.	146
35	Correlation coefficient between heavy metal concentrations in seaweed and seawater physicochemical parameters.	147
36	Statistical analysis of the metal bioaccumulation data (Kruskal-Wallis one way analysis of variance).	148
37	Concentration (means \pm S.D) in $\mu\text{g g}^{-1}$ dry weight of heavy metals in <i>E. intestinalis</i>	150
38	Correlations between concentrations of paired metals in <i>Cladophoropsis zollingeri</i> .	150
39	Correlations between concentrations of paired metals in <i>Chaetomorpha aerea</i> .	150
40	Correlations between concentrations of paired metals in <i>Enteromorpha intestinalia</i> .	151
41	List of brown algae sampled from different sites.	151
42	Results obtained from the multiple regression analysis of the metal contents in Phaeodophyta species in relationship with the metal contamination in sediment and seawater.	152
43	Correlations between concentrations of paired metals in <i>Padina</i> sp.	155
44	Correlations between concentrations of paired metals in <i>C. myrica</i> .	155
45	Correlations between concentrations of paired metals in <i>C. sinousa</i> .	155
46	Correlations between concentrations of paired metals in <i>Sargassum</i> sp.	157
47	Description of red algae sampled in different sites.	158
48	Results obtained from the multiple regression analysis of the metal contents in Rhodophyta species in relationship with the metal contamination in sediment and seawater.	159
49	Correlations between concentrations of paired metals in <i>L. obtusa</i> .	160
50	Correlations between concentrations of paired metals in <i>J. rubens</i> .	162
51	Correlations between concentrations of paired metals in <i>C. parvula</i> .	162

52	Results of statistical analysis based on estimates of heavy metal concentrations in the three classes of seaweed and seawater around the Kish Island.	164
53	Results of statistical analysis based on estimates of concentrations factors in three classes of seaweeds around the Kish Island.	166
54	Concentrations factors of different classes of seaweeds during sampling periods around the Kish Island.	166
55	Heavy metals concentrations (means \pm S.D.) in different parts of <i>Cystoseira myrica</i> and <i>Sargassum ilicifolium</i> .	167
56	Heavy metals concentrations (mean \pm S.D.) in $\mu\text{g g}^{-1}$ dry weight in three classes of seaweeds during different periods around the Kish Island.	171
57	Mean of MPI amounts in sediment, seawater and seaweeds in Kish Island	181
58	Effects of Cd on the <i>C. myrica</i> after 14 days (means \pm S.D.).	184
59	Effects of Cu on the <i>C. myrica</i> after 14 days (means \pm S.D.).	187
60	Effects of Ni on the <i>C. myrica</i> after 14 days (means \pm S.D.).	188
61	Effects of Pb on the <i>C. myrica</i> after 14 days (means \pm S.D.).	190
62	Effects of V on the <i>C. myrica</i> after 14 days (means \pm S.D.).	192
63	Effects of Zn on the <i>C. myrica</i> after 14 days (means \pm S.D.).	194
64	Effects of Cd on photosynthesis, respiration and chlorophyll contents in <i>S. ilicifolium</i> after 96 hrs (means \pm S.D.).	196
65	Effects of Cu on photosynthesis, respiration and chlorophyll contents in <i>S. ilicifolium</i> after 96 hrs (means \pm S.D.).	199
66	Effects of Ni on photosynthesis, respiration and chlorophyll contents in <i>S. ilicifolium</i> after 96 hrs (means \pm S.D.).	201
67	Effects of Pb on photosynthesis, respiration and chlorophyll contents in <i>S. ilicifolium</i> after 96 hrs (means \pm S.D.).	203
68	Effects of V on photosynthesis, respiration and chlorophyll contents in <i>S. ilicifolium</i> after 96 hrs (means \pm S.D.).	205

69	Effects of Zn on photosynthesis, respiration and chlorophyll contents in <i>S. ilicifolium</i> after 96 hrs (means \pm S.D.).	207
70	The estimated of the EC ₅₀ values for single metals using the chlorophyll reduction.	207
71	Exponential decay non-linear regression to <i>Cystoseira myrica</i> and <i>Sargassum ilicifolium</i> in relationship with metal exposure at different concentrations.	209
72	Total extent and efficiency of metals bioaccumulation (concentration factor) (means \pm S.D.) during two weeks by <i>C. myrica</i> .	216
73	Metals accumulation (means \pm S.D.) during two weeks toxicity test by <i>C. myrica</i> .	217
74	Total extent and efficiency of metals bioaccumulation (concentration factor) (means \pm S.D.) during 96 hrs by <i>S. ilicifolium</i> .	219
75	Metals accumulation (means \pm S.D.) during 96 hrs toxicity test by <i>S. ilicifolium</i> .	220
76	Heavy metal concentrations (mg l ⁻¹) in different parts of <i>S. ilicifolium</i> (means \pm S.D.).	223
77	Efficiency of metals bioaccumulation (concentration factor) during 12 hrs incubation in 0.1mg l ⁻¹ solution to light and dark periods (means \pm S.D.).	223
78	Correlation between initial metal concentration in seawater and metal accumulated by <i>C. myrica</i> .	227
79	Correlation coefficient between initial metal concentration in seawater and metal accumulated by <i>S. ilicifolium</i> .	228
80	Characteristics of the sampling sites.	231

LIST OF FIGURES

Figures		Page
1	Major river systems, depth contours (in meters), relation to Africa (Insert A), and tectonic plates (Insert B) of the Persian Gulf.	2
2	Relationship between contamination and pollution.	54
3	Study area map showing the study area locations.	61
4	Axis morphology of primary branches of <i>Sargassum</i> .	80
5	Zonation of shoreline.	102
6	Dendrogram using average linkage to different groups of seaweeds between sites	102
7	Seasonal biomass (g wet wt m ⁻²) of seaweeds at different locations in the Kish Island.	114
8	The mean seasonal biomass of 3 groups of seaweeds in the Kish Island.	115
9	Relative biomass of 3 groups of seaweeds in the Kish Island.	115
10	Variations of air temperature and total algal biomass (means \pm S.D.) during different seasons.	122
11	Variations of nutrients with total algal biomass (means \pm S.D.) in different seasons.	122
12	The relation between total algal biomass and total bright sunshine (means \pm S.D.) during each season.	122
13	The relation between total algal biomass and wind speed (means \pm S.D.) during each season.	123
14	Circulation of surface water in the Persian Gulf.	131
15	Seasonal concentrations (means \pm S.D.) of heavy metals in $\mu\text{g l}^{-1}$ during different seasons in coastal waters of Kish Island.	132
16	Dendrogram showing classification of stations based on capacity of metals uptake by seaweeds.	156
17	Dendrogram showing classification of stations based on capacity of metals uptake by seaweeds.	160

18	Metals concentration (means \pm S.D.) in <i>Cystoseira myrica</i> . A: Different age B: Different parts	169
19	Metals concentration (means \pm S.D.) in <i>Sargassum</i> sp.	169
20	Seasonal variation of metal concentration ($\mu\text{g g}^{-1}$ dry weight) in three class of seaweeds from Kish Island.	174
21	MPI values in seaweeds at each sampling site.	181
22	MPI values in sediment and seawater during different periods.	182
23	MPI values in three classes of seaweeds during different sampling periods.	182
24	Effect of different Cd concentrations (mg l^{-1}) on the growth rate (\pm S.D.) of <i>C. myrica</i> .	185
25	Variation in chlorophyll <i>a</i> (\pm S.D.) during Cd toxicity test on <i>C.</i> <i>myrica</i> .	185
26	Effect of different Cu concentrations (mg l^{-1}) on the growth rate (\pm S.D.) of <i>C. myrica</i> .	187
27	Variation in chlorophyll <i>a</i> (\pm S.D.) during Cu toxicity test on <i>C.</i> <i>myrica</i> .	187
28	Effect of different Ni concentrations (mg l^{-1}) on the growth rate (\pm S.D.) of <i>C. myrica</i> .	189
29	Variation in chlorophyll <i>a</i> (\pm S.D.) during Ni toxicity test on <i>C.</i> <i>myrica</i> .	189
30	Effect of different Pb concentrations (mg l^{-1}) on the growth rate (\pm S.D.) of <i>C. myrica</i> .	191
31	Variation in chlorophyll <i>a</i> (\pm S.D.) during Pb toxicity test on <i>C.</i> <i>myrica</i> .	191
32	Effect of different V concentrations (mg l^{-1}) on the growth rate (\pm S.D.) of <i>C. myrica</i> .	192
33	Variation in chlorophyll <i>a</i> (\pm S.D.) during V toxicity test on <i>C.</i> <i>myrica</i> .	193
34	Effect of different Zn concentrations (mg l^{-1}) on the growth rate (\pm S.D.) of <i>C. myrica</i> .	194

35	Variation in chlorophyll <i>a</i> (\pm S.D.) during Zn toxicity test on <i>C. myrica</i> .	194
36	Variation of chlorophyll <i>a</i> (\pm S.D.) during Cd toxicity test on <i>S. ilicifolium</i> .	197
37	Variation of chlorophyll <i>a</i> (\pm S.D.) during Cu toxicity test on <i>S. ilicifolium</i> .	199
38	Variation of chlorophyll <i>a</i> (\pm S.D.) during Ni toxicity test on <i>S. ilicifolium</i> .	201
39	Variation of chlorophyll <i>a</i> (\pm S.D.) during Pb toxicity test on <i>S. ilicifolium</i> .	203
40	Variation of chlorophyll <i>a</i> (\pm S.D.) during V toxicity test on <i>S. ilicifolium</i> .	205
41	Variation of chlorophyll <i>a</i> (\pm S.D.) during Zn toxicity test on <i>S. ilicifolium</i> .	207
42	Exponential decay non-linear regression of the different metal concentrations on the <i>Cystoseira myrica</i> .	210
43	Exponential decay non-linear regression of the different metal concentrations on the <i>Sargassum ilicifolium</i> .	212
44	Comparison of heavy metals accumulated (means \pm S.D.) in different metals concentrations by <i>C. myrica</i> .	218
45	Comparison of heavy metals accumulated (means \pm S.D.) in different metals concentrations by <i>S. ilicifolium</i> .	222
46	Heavy metal accumulation (means \pm S.D.) in three different parts of <i>S. ilicifolium</i> during light period.	224
47	Heavy metal accumulation (means \pm S.D.) in three different parts of <i>S. ilicifolium</i> sp. during dark period.	224
48	Relation between metal concentration with the rate of metal uptake by <i>C. myrica</i> .	229
49	Relation between metal concentration with the rate of metal uptake in <i>S. ilicifolium</i> .	229

LIST OF PLATES

Plates		Page
1	Study area (site 1) during low tide	63
2	Study area (site 2) during low tide	63
3	Study area (site 3) during low tide	64
4	Study area (site 4) during low tide	64
5	Study area (site 5) during low tide	65
6	Desalination Plant Site	65
7	A 50 × 50 cm square metal frame use for sampling	67
8	<i>Enteromorpha intestinalis</i>	88
9	<i>Chaetomorpha aerea</i>	89
10	<i>Cladophora sericioides</i>	91
11	<i>Cladophoropsis zollingeri</i>	91
12	<i>Dictyosphaeria cavernosa</i>	91
13	<i>Caulerpa racemosa</i>	91
14	<i>Colpomenia sinousa</i>	94
15	<i>Padina</i> sp.	94
16	<i>Cystoseira myrica</i>	94
17	<i>Sargassum latifolium</i>	94
18	<i>Sargassum polyceratium</i>	94
19	<i>Sargassum vulgar</i>	95
20	<i>Sargassum polycystum</i>	95
21	<i>Turbinaria conoides</i>	96
22	<i>Liagora ceranoides</i>	97
23	<i>Jania rubens</i>	97

24	<i>Gelidium</i> sp.	98
25	<i>Gracilaria foliifera</i>	98
26	<i>Sarconema filiiforma</i>	99
27	<i>Champia parvula</i>	100
28	<i>Acanthophora spicifera</i>	100
29	<i>Laurencia obtusa</i>	101
30	<i>Laurencia</i> sp.	101
31	Section of <i>Cystoseira myrica</i> (a) before and (b) after exposure to 1 mg l ⁻¹ of Cu ⁺²	186

LIST OF ABBREVIATIONS

A.A.S.	=	Atomic Absorption Spectrophotometer
ANOVA	=	Analysis of Variance
APDC	=	Ammonium Pyrrolidine Dithiocarbamate
COMAS	=	Center for Oceanography and Mariculture Studies
CV	=	Coefficient of Variations
D.O.	=	Dissolved Oxygen
D W	=	Dry weight
EC ₅₀	=	Effective Concentration resulting 50% mortality of test Organism
EN	=	Electronegativity
g wet wt m ⁻²	=	Grams Wet Weight per square meter
I.C.P-A.E.S	=	Inductively Coupled Plasma-Atomic Emission Spectrophotometer
mg l ⁻¹	=	milligram per liter
MIBK	=	Methyl Iso-Butyl Ketone
MPI	=	Metal Pollution Index
NED	=	Naphthylethylenediamine dihydrochloride
ROPME	=	Regional Organization for the Protection of the Marine Environment
St.	=	Site
µg l ⁻¹	=	microgram per liter
mg g ⁻¹	=	milligram per gram
µg g ⁻¹	=	microgram per gram
ppm	=	Part per million
ppb	=	Part per billion

<i>P</i>	=	Probability
P.S.	=	Photosynthesis System
TES	=	Titrisol Eisen Standard
Vs	=	Versus
UAE	=	United Arab Emirates
Wet wt.	=	Wet weight